


Update of SeaQuest Data Analysis on $\bar{d}/\bar{u}$ Ratio of Nucleons



Kei Nagai
Academia Sinica

26th November, 2018



The 2nd Workshop on Parton Distribution Functions @ Nangang, Taipei, Taiwan



Introduction



Antiquark Flavor Asymmetry

- Gluon splitting: Flavor Independent $\bar{u} = \bar{d}$
- Gottfried sum

$$S_G \equiv \int_0^1 \frac{dx}{x} [F_2^p(x) - F_2^n(x)] = \frac{1}{3} + \frac{2}{3} \left(\int_0^1 \bar{u}_p(x) dx - \int_0^1 \bar{d}_p(x) dx \right)$$

x : Bjorken $x = P_{\text{parton}}/P_{\text{proton}}$ @ high-energy


Assumption: PDFs in neutron and proton are isospin symmetric:

$$\int_0^1 u_p(x) dx = \int_0^1 d_n(x) dx, \dots$$

- NMC experiment at CERN (1990, DIS)

$$S_G = 0.235 \pm 0.026 < 1/3$$

$$\rightarrow \int_0^1 \bar{d}(x) dx > \int_0^1 \bar{u}(x) dx$$






Antiquark Flavor Asymmetry

- NA51 Experiment (Drell-Yan) @ CERN
 - **Significant Flavor Asymmetry**
 $\bar{d}/\bar{u} = 1.96 @ x = 0.18$
- E866 Experiment (Drell-Yan) @ Fermilab
 - **Significant Flavor Asymmetry**
 $\bar{d}/\bar{u} \sim 1.7 @ x \sim 0.2$
 - $\bar{d}/\bar{u} < 1.0 @ x \sim 0.3?$
with large stat. uncertainty
No theories can reproduce

$$x : \text{Bjorken } x = \frac{P_{\text{parton}}}{P_{\text{proton}}}$$

(@ high energy)





Models for $\bar{d}/\bar{u}$

- Pauli Blocking

- $g \rightarrow u\bar{u}$ is suppressed compared to $g \rightarrow d\bar{d}$ ($p = uud$)
- Only few % effect [NPB149, 497 (1979)]

- Statistical model [NPA948, 63 (2016)]

- Fermi (quarks) and Bose (gluons) statistics

- Meson cloud model [PRD58, 092004 (1998)]


- $|p\rangle = |p_0\rangle + \alpha|N\pi^+\rangle + \beta|\Delta\pi^-\rangle + \gamma|\Lambda K\rangle + \dots$

★ $N\pi^+ = (udd)(u\bar{d})$

★ $\Delta\pi^- = (uuu)(d\bar{u})$

★ $\alpha > \beta$

→ $\bar{d} > \bar{u}$



- etc...

SeaQuest will provide the new data points ($0.1 < x < 0.45$)
and it is important to understand the structure of the proton!



SeaQuest Experiment



SeaQuest Experiment

Drell-Yan experiment

- Performed at Fermilab (Illinois, US)
 - Main Injector
 - 120 GeV ($\sqrt{s} \sim 15$ GeV) proton beam
 - 5 seconds of beam is provided every 60 seconds (other 55 seconds for neutrino experiments)
 - 53 MHz beam bunch, $\sim 40k$ protons in a bunch
- Topics
 - **Antiquark Flavor Asymmetry**
 - Nuclear dependence
 - Angular Distribution
 - Dark Photon Search





Collaboration List

Abilene Christian University

A. Boles, B. Bowen, K. Bowling, A.T. Brown, A.W. Brown,
E. Carlisle, P. Carstens, R. Castillo, M. Crowder,
M. Daugherty, B. Edlund, T. Hague, L.D. Isenhower,
N. Kitts, S. Marshall, L. Medlock, A. Miller, B. Miller,
L. Selensky, T. Sipos, R. Towell, S. Watson, Z. Xi

Academia Sinica

T.-H. Chang, W.-C. Chang, K. Nagai, S.-H. Shiu, D.-S. Su,
M. Quaresma

Argonne National Laboratory

J. Arrington, C. Durandet, J. Ferreri, [D.F. Geesaman*](#),
R.J. Holt, M.M. Medeiros, [P.E. Reimer*](#), B.G. Tice, Z. Ye

University of Colorado

E. Erdos, J. Katich, E. Kinney, P.-J. Lin, B. McDonald

Fermi National Accelerator Laboratory

C. Brown, D. Christian, J.-Y. Wu

University of Illinois

B. Dannowitz, A. Chen, M. Diefenthaler, B. Kerns, J. Kras,
H. Li, D. Liu, N.C.R. Makins, R.E. McClelland, J.-C. Peng,
S. Prasad, M.H. Teo, M. Witek, Y. Yin

KEK

S. Sawada

Los Alamos National Laboratory

G.T. Garvey, L. Guo, X. Jiang, A. Klein, D. Kleinjan,
M. Leitch, H. Liu, K. Liu, M. Liu, P. McGaughey, A. Puckett

Mississippi State University

L. El Fassi

University of Maryland

S. Bastola, E. Beise, K. Nakahara

University of Michigan

C.A. Aidala, C. Ayuso, C. Culklin, C. Dutta, V.R. Logins,
W. Lorenzon, I.A. Mooney, D. Morton, B. Nadim,
B.J. Ramson, R. Raymond, J.G. Rubin, T. Sawada, M. Scott,
M. Stewart, M.R. Wood

National Kaohsiung Normal University

R. Guo, S.-Y. Wang

RIKEN

Y. Goto

Rutgers University

R. Gilman, R. Ransome, A.S. Tadepalli, Y. Zhang

Tokyo Institute of Technology

S. Miyasaka, K. Nakano, S. Obata, F. Sanftl, T.-A. Shibata,
S. Takeuchi

Yamagata University

Y. Kudo, Y. Miyachi, S. Nara

* Co-Spokespersons






Drell-Yan Process

- Cross section ($p + p$, Leading order)


$$\frac{d^2\sigma}{dx_{\text{target}}dx_{\text{beam}}} = \frac{4\pi\alpha^2}{9x_{\text{target}}x_{\text{beam}}} \frac{1}{s} \sum_i e_i^2 [q_i(x_{\text{beam}})\bar{q}_i(x_{\text{target}}) + \bar{q}_i(x_{\text{beam}})q_i(x_{\text{target}})]$$

- An antiquark is always involved
- $\bar{q}(x_{\text{beam}})q(x_{\text{target}}) \sim 0$ @ $x_{\text{beam}} \gg x_{\text{target}}$
 - Access **antiquarks in target proton** and **quarks in beam proton**
- Final state dimuons are measured






SeaQuest Spectrometer




- Targets: LH_2 , LD_2 , C, Fe, W
- Hadron Absorbers (stop beam, muon identification)
- Magnets (focussing, momentum determination)
- 4 tracking stations, consist of
 - Hodoscopes
 - Drift Chambers (St. 1-3) or Prop. Tubes (St. 4)



- Mass distribution fitted with estimated components
- Well fitted:
Detectors & tracking tool work as expected
- Drell-Yan can be selected with
mass $> 4.2 \text{ GeV}/c^2$



Timeline



Year	Month	
2011	08	Finish spectrometer construction
2012	03-04	Commissioning data taking (Run I)
	05-	Detector upgrade
2013	11-	Phys. data taking (Run II)
2014	-09	
2015	11-	Phys. data taking (Run III)
	-07	Phys. data taking (Run IV)
2016	10-	
2016	-08	Phys. data taking (Run V)
	11-	
2017	-07	

- Finished data taking (2017.07)
- Recorded protons on targets: 1.4×10^{18}
- Status in FY2017
 - 0.3×10^{18}
 - Wider St. 1 chamber:
 - 40% more effective for large x (~ 0.4)



Data Analysis



Extract $\bar{d}/\bar{u}$

Basic Idea


$$\frac{\sigma_{pd}(x)}{2\sigma_{pp}(x)} \approx \frac{1}{2} \left[1 + \frac{\bar{d}(x)}{\bar{u}(x)} \right] \quad \left| \quad \begin{array}{l} \text{Drell-Yan cross section ratio is proportional to } \bar{d}/\bar{u} \\ \text{with } x_{\text{beam}} \gg x_{\text{target}} \end{array} \right.$$

- Cross section ratio: $\frac{\sigma_{pd}}{2\sigma_{pp}} = \frac{1}{2} \left(\frac{N_D \cdot C_D}{P_D} \right) \bigg/ \left(\frac{N_H \cdot C_H}{P_H} \right)$
 - Number of dimuons (N)
 - Background and reconstruction efficiency corrections (C)
 - Normalization with number of nucleons in beam and target (P)
- Convert $\sigma_{pd}/2\sigma_{pp}$ to $\bar{d}/\bar{u}$
 - $\frac{d^2\sigma}{dx_{\text{target}}dx_{\text{beam}}} = \frac{4\pi\alpha^2}{9x_{\text{target}}x_{\text{beam}}} \frac{1}{s} \sum_i e_i^2 [q_i(x_{\text{beam}})\bar{q}_i(x_{\text{target}}) + \bar{q}_i(x_{\text{beam}})q_i(x_{\text{target}})]$
LO Drell-Yan cross section is used for extracting $\bar{d}/\bar{u}$



Cross Section Ratio

Preliminary result



Systematics

- H contamination in LD₂
- background
- hit-rate dependence of reconstruction efficiency


Statistics
Cross section ratio of σ^{pd} to σ^{pp}

- LH₂: 17951 dimuons,
LD₂: 20284 dimuons

- Cross section ratio at $0.1 < x < 0.58$ is obtained
- $\sigma_{pd}/2\sigma_{pp} > 1.0$ @ $0.10 < x_2 < 0.45$
 $\sigma_{pd}/2\sigma_{pp} = 1.0$ @ $0.45 < x_2 < 0.58$



$\bar{d}/\bar{u}$ Preliminary Result



Systematic uncertainty

- H contamination in LD₂
- background
- hit-rate dependence of reconstruction efficiency
- uncertainty from CT10 PDF (cross section ratio $\rightarrow \bar{d}/\bar{u}$)

Note: Nuclear corrections for deuterium have not yet been applied.

SeaQuest Preliminary Result (LO)

- $\bar{d}/\bar{u} > 1.0$ @ $0.10 < x < 0.45$
- $\bar{d}/\bar{u} = 1.0$ @ $0.45 < x < 0.58$
within stats. error

Comparison with NA51, E866

- $0.1 < x < 0.24$: well consistent
- $x > 0.24$: SeaQuest $>$ E866 !?
 - Difference of Q^2 ? – No effect
 - Difference of PDF sets? – No effect




Toward Final Results



Difficulties in Current Analysis?

- Reconstruction efficiency effect
 - Reconstruction efficiency decreases as intensity increases (rate-dependent)



- Combinatorial background
 - Example: Single muon coming from π + single muon from Drell-Yan
 - Estimated with event mixing method
 - ★ Mix muons from different events
 - Background would be rate-dependent


How to remove these effects?

- Basic Idea: These effects should vanish at “zero” intensity



Extrapolation Method

- Cross-section ratio as a function of intensity
- Extrapolate the data to “Zero” intensity
- **Intercept value is “correct” cross-section ratio**
 - No reconstruction efficiency effect
 - No combinatorial background



Difficulties of this method?

- Fitting shape?
- Validity of the method?
- How to extract $\bar{d}/\bar{u}$?

Investigation is in progress...



Summary

- SeaQuest aims to investigate the structure of the proton using Drell–Yan process.
- Drell–Yan process is sensitive to the antiquark distributions. Suitable for the investigation of the antiquark flavor asymmetry.
- Antiquark flavor asymmetry is important to understand the structure of the proton.
- SeaQuest finished the data taking (2012-2017).
- The preliminary results from FY2015 were released.
 - $\bar{d}/\bar{u} > 1.0$ @ $0.10 < x < 0.45$,
 - $\bar{d}/\bar{u} = 1.0$ @ $0.45 < x < 0.58$
 - Further investigation of the discrepancy between E866.
- Works for final results are in progress.
 - New method: Extrapolation method
 - Resolving the difficulties of this method. Final results will be released soon!



Backup