

Towards ideal topological materials: Comprehensive database searches using symmetry indicators

Xiangang Wan

(万贤纲)


Department of Physics, Nanjing University, China

The 17th Workshop on First-principles Computational
Materials Physics

2019年6月25日 @Tsinghua University, Taiwan

Cooperation

Feng Tang; *Nanjing University*



Hoi Chun Po & Ashvin Vishwanath; *Harvard University*



*Tang, Po, Vishwanath, **Wan***, Nature Physics 15, 470 (2019)*

*Tang, Po, Vishwanath, **Wan***, Science Advances 5, eaau8725 (2019)*

*Tang, Po, Vishwanath, **Wan***, Nature 566, 486 (2019)*

Content


Symmetry Indicators

Topological Materials

拓扑

➤ 朗道相变理论 → 对称性、序参量

➤ 1980



Quantized Hall Conductance in a Two-Dimensional Periodic Potential

D. J. Thouless, M. Kohmoto,^(a) M. P. Nightingale, and M. den Nijs

Department of Physics, University of Washington, Seattle, Washington 98195

(Received 30 April 1982)

The Hall conductance of a two-dimensional electron gas has been studied in a uniform magnetic field and a periodic substrate potential U . The Kubo formula is written in a form that makes apparent the quantization when the Fermi energy lies in a gap. Explicit expressions have been obtained for the Hall conductance for both large and small $U/\hbar\omega_c$.

$$\sigma_H = \frac{ie^2}{A_0 \hbar} \sum_{\epsilon_\alpha < E_F} \sum_{\epsilon_\beta > E_F} \frac{(\partial \hat{H} / \partial k_1)_{\alpha\beta} (\partial \hat{H} / \partial k_2)_{\beta\alpha} - (\partial \hat{H} / \partial k_2)_{\alpha\beta} (\partial \hat{H} / \partial k_1)_{\beta\alpha}}{(\epsilon_\alpha - \epsilon_\beta)^2},$$

Chern Number


- 2D BZ是一个torus，它组成的曲面包含的monopole的个数只能是整数

$$\sigma_H = \frac{ie^2}{A_0 \hbar} \sum_{\epsilon_\alpha < E_F} \sum_{\epsilon_\beta > E_F} \frac{(\partial \hat{H} / \partial k_1)_{\alpha\beta} (\partial \hat{H} / \partial k_2)_{\beta\alpha} - (\partial \hat{H} / \partial k_2)_{\alpha\beta} (\partial \hat{H} / \partial k_1)_{\beta\alpha}}{(\epsilon_\alpha - \epsilon_\beta)^2}$$

$$\Omega_{\mu\nu}^n(\mathbf{R}) = i \sum_{n' \neq n} \frac{\langle n | \partial H / \partial R^\mu | n' \rangle \langle n' | \partial H / \partial R^\nu | n \rangle - (\nu \leftrightarrow \mu)}{(\epsilon_n - \epsilon_{n'})^2}.$$

$$\sigma_{xy} = \frac{e^2}{\hbar} \int_{\text{BZ}} \frac{d^2 k}{(2\pi)^2} \Omega_{k_x k_y} = n \frac{e^2}{h}.$$

Chern Number



Z2 Topological insulator

PRL **95**, 226801 (2005)

PHYSICAL REVIEW LETTERS

week ending
25 NOVEMBER 2005

Quantum Spin Hall Effect in Graphene

C. L. Kane and E. J. Mele


PRL **95**, 146802 (2005)

PHYSICAL REVIEW LETTERS

30

Z₂ Topological Order and the Quantum Spin Hall Effect

C. L. Kane and E. J. Mele



Z₂拓扑不变量

时间反演对称

SOC


Topological invariants in energy band

- **Chern number** in T -symmetry breaking system
- **Z_2 topological invariant** in T -symmetry invariant system

Topological Insulator

- Bulk insulator
- Topological protected surface state
- Mass term in Dirac Equation change Sign

$$H_t = \alpha \cdot \mathbf{p} + \beta m$$



Bulk-boundary correspondence




FIG. 3. (Color online) Electronic dispersion between two boundary Kramers degenerate points $\Gamma_a=0$ and $\Gamma_b=\pi/a$. In (a) the number of surface states crossing the Fermi energy E_F is even, whereas in (b) it is odd. An odd number of crossings leads to topologically protected metallic boundary states.

Weak topological insulator

Z_2 invariants $(\gamma_0; \gamma_1, \gamma_2, \gamma_3)$

Strong topological insulator

$$\mathcal{H}_{\text{surface}} = -i\hbar v_F \vec{\sigma} \cdot \vec{\nabla},$$

Justify 3D TI

- ① Surface State Electron Structure
- ② Adiabatic Continue Band Transformation
- ③ Band inversion
- ④ Calculate Z_2

Calculate Z_2

➤ With inversion symmetry

Fu-Kane theorem (*Fu & Kane, 2007*)

➤ Without inversion symmetry

directly calculate Z_2

(*Feng, Wen, Zhou, Xiao & Yao (2012)*)

$$Z_2 = \frac{1}{2\pi} \left[\oint_{\partial\mathcal{B}^+} dk \cdot \mathcal{A}(k) - \int_{\mathcal{B}^+} d^2k \mathcal{F}(k) \right] \text{ mod } 2,$$

where $\mathcal{A}(k)$ and $\mathcal{F}(k)$ are the Berry connection and Berry curvature, respectively,

$$\mathcal{A}(k) = i \sum_n \langle u_n(k) | \nabla_k u_n(k) \rangle \quad \mathcal{F}(k) = \nabla_k \times \mathcal{A}(k) \Big|_z .$$

周期表

PHYSICAL REVIEW B 78, 195125 (2008)

Classification of topological insulators and superconductors in three spatial dimensions

Andreas P. Schnyder,¹ Shinsei Ryu,¹ Akira Furusaki,² and Andreas W. W. Ludwig³

		TRS	PHS	SLS	$d=1$	$d=2$	$d=3$
Standard (Wigner-Dyson)	A (unitary)	0	0	0	-	\mathbb{Z}	-
	AI (orthogonal)	+1	0	0	-	-	-
	AII (symplectic)	-1	0	0	-	\mathbb{Z}_2	\mathbb{Z}_2
Chiral (sublattice)	AIII (chiral unitary)	0	0	1	\mathbb{Z}	-	\mathbb{Z}
	BDI (chiral orthogonal)	+1	+1	1	\mathbb{Z}	-	-
	CII (chiral symplectic)	-1	-1	1	\mathbb{Z}	-	\mathbb{Z}_2
BdG	D	0	+1	0	\mathbb{Z}_2	\mathbb{Z}	-
	C	0	-1	0	-	\mathbb{Z}	-
	DIII	-1	+1	1	\mathbb{Z}_2	\mathbb{Z}_2	\mathbb{Z}
	CI	+1	-1	1	-	-	\mathbb{Z}

TABLE I. Ten symmetry classes of single-particle Hamiltonians classified in terms of the presence or absence of time-reversal symmetry (TRS) and particle-hole symmetry (PHS), as well as “sublattice” (or “chiral”) symmetry (SLS) (Refs. 37 and 38). In the table, the absence of symmetries is denoted by “0.” The presence of these symmetries is denoted by either “+1” or “-1,” depending on whether the (antiunitary) operator implementing the symmetry at the level of the single-particle Hamiltonian squares to “+1” or “-1” (see text). [The index ± 1 equals η_c in Eq. (1b); here $\epsilon_c = +1$ and -1 for TRS and PHS, respectively.] For the first six entries of the table (which can be realized in nonsuperconducting systems), TRS = +1 when the SU(2) spin is an integer [called TRS (even) in the text] and TRS = -1 when it is a half-integer [called TRS (odd) in the text]. For the last four entries, the superconductor “Bogoliubov–de Gennes” (BdG) symmetry classes D, C, DIII, and CI, the Hamiltonian preserves SU(2) spin-1/2 rotation symmetry when PHS = -1 [called PHS (singlet) in the text], while it does not preserve SU(2) when PHS = +1 [called PHS (triplet) in the text]. The last three columns list all topologically non-trivial quantum ground states as a function of symmetry class and spatial dimension. The symbols \mathbb{Z} and \mathbb{Z}_2 indicate whether the space of quantum ground states is partitioned into topological sectors labeled by an integer or a \mathbb{Z}_2 quantity, respectively.

Topological Crystalline Insulators

Spatial Symmetries

PRL **106**, 106802 (2011)

PHYSICAL REVIEW LETTERS

Topological Crystalline Insulators

Liang Fu

ARTICLES

PUBLISHED ONLINE: 16 DECEMBER 2012 | DOI:10.1038/NPHYS2513

nature
physics

The space group classification of topological band-insulators

Robert-Jan Slager¹, Andrej Mesaros², Vladimir Juričić^{1*} and Jan Zaanen¹

Y. Ando and L. Fu, *Annual Review of Condensed Matter Physics* **6**, 361 (2015).

Topological Crystalline Insulators

➤ Mirror Chern Insulator

Ando and Fu, Annu. Rev. Condens. Matter Phys. 6, 361 (2015)

➤ Hourglass Fermions

Wang, Alexandradinata, Cava, Bernevig, Nature 532, 189 (2016).

➤ Wallpaper Fermions

Wieder et al., Science 361, 246 (2018)

➤ Higher-order topological insulators

Fang&Fu; Schindler et al., Langbehn et al., Song, Fang&Fang; Benalcazar et al., (2017)

...


Higher-Order Topological Insulators

arXiv:1708.03636

Frank Schindler,¹ Ashley M. Cook,¹ Maia G. Vergniory,^{2,3} Zhijun Wang,⁴
Stuart S. P. Parkin,⁵ B. Andrei Bernevig,^{4,2,6} and Titus Neupert¹

bulk $(d) \rightarrow d-1, d-2, d-3$

Second order



Topological Insulators Turn a Corner

Theorists have discovered topological insulators that are insulating in their interior and on their surfaces but have conducting channels at corners or along edges.

by Siddharth A. Parameswaran* and Yuan Wang†

- [2] W. A. Benalcazar, B. A. Bernevig, and T. L. Hughes, “Quantized Electric Multipole Insulators,” *Science* **357**, 61 (2017).
- [3] W. A. Benalcazar, B. A. Bernevig, and T. L. Hughes, “Electric Multipole Moments, Topological Multipole Moment Pumping, and Chiral Hinge States in Crystalline Insulators,” *Phys. Rev. B* **96**, 245115 (2017).
- [4] Z. Song, Z. Fang, and C. Fang, “ $(d - 2)$ -dimensional edge states of rotation symmetry protected topological states,” *Phys. Rev. Lett.* **119**, 246402 (2017).
- [5] J. Langbehn, Y. Peng, L. Trifunovic, F. von Oppen, and P. W. Brouwer, “Reflection-symmetric second-order topological insulators and superconductors,” *Phys. Rev. Lett.* **119**, 246401 (2017).




Figure 1: Usually, 3D topological insulators conduct via gapless states on their 2D surfaces but are insulating in their bulk (left). Recently proposed second- and third-order 3D TIs have gapless states on their 1D hinges (middle) or 0D corners (right), respectively, and they constitute a new class of topological phases of matter. (APS/Alan Stonebraker)


11 December 2017 *Physics* **10**, 132

Dirac Fermion in Condensed Matter

Massless Dirac Fermion

K-point at BZ of Graphene (1947)

2004, 2005



Two-dimensional gas of massless Dirac fermions in graphene

Vol 438|10 November 2005|doi:10.1038/nature04233

K. S. Novoselov¹, A. K. Geim¹, S. V. Morozov², D. Jiang¹, M. I. Katsnelson³, I. V. Grigorieva¹, S. V. Dubonos²
& A. A. Firsov²

Vol 438|10 November 2005|doi:10.1038/nature04235

Experimental observation of the quantum Hall effect and Berry's phase in graphene

Yuanbo Zhang¹, Yan-Wen Tan¹, Horst L. Stormer^{1,2} & Philip Kim¹

Dirac Fermion in CM


Massless Dirac Fermion 3D

symmetry protected degeneracies

Dirac Semimetal in Three Dimensions

PRL **108**, 140405 (2012)

S. M. Young,¹ S. Zaheer,² J. C. Y. Teo,^{2,*} C. L. Kane,² E. J. Mele,² and A. M. Rappe¹



- **Four dimensional irreducible representations at high symmetry points of BZ**
- **Linearly in all directions around these points**

Dirac Fermion in CM

Dirac semimetal and topological phase transitions in $A_3\text{Bi}$ ($A = \text{Na, K, Rb}$)

Zhijun Wang,¹ Yan Sun,² Xing-Qiu Chen,² Cesare Franchini,² Gang Xu,¹ Hongming Weng,^{1,*} Xi Dai,¹ and Zhong Fang^{1,†}


PHYSICAL REVIEW B **85**, 195320 (2012)

Three-dimensional Dirac semimetal and quantum transport in Cd_3As_2

Zhijun Wang, Hongming Weng,^{*} Quansheng Wu, Xi Dai, and Zhong Fang[†]

PHYSICAL REVIEW B **88**, 125427 (2013)

- **Two bands along high symmetry lines of BZ belong to different two dimensional irreducible representations →**




Dirac Fermion in CM

Discovery of a Three-Dimensional Topological Dirac Semimetal, Na_3Bi


Z. K. Liu,^{1*} B. Zhou,^{2,3*} Y. Zhang,³ Z. J. Wang,⁴ H. M. Weng,^{4,5} D. Prabhakaran,² S.-K. Mo,³
Z. X. Shen,¹ Z. Fang,^{4,5} X. Dai,^{4,5} Z. Hussain,³ Y. L. Chen^{2,6†}

21 FEBRUARY 2014 VOL 343 SCIENCE



A stable three-dimensional topological Dirac semimetal Cd_3As_2

Z. K. Liu^{1†}, J. Jiang^{2,3†}, B. Zhou^{2,4†}, Z. J. Wang^{5†}, Y. Zhang^{1,4}, H. M. Weng⁵, D. Prabhakaran², S.-K. Mo⁴,
H. Peng², P. Dudin⁶, T. Kim⁶, M. Hoesch⁶, Z. Fang⁵, X. Dai⁵, Z. X. Shen¹, D. L. Feng³, Z. Hussain⁴
and Y. L. Chen^{1,2,4,6*}




PUBLISHED ONLINE: 25 MAY 2014 | DOI: 10.1038/NMAT3990

Correlation \rightarrow band structure evolution


$\text{Y}_2\text{Ir}_2\text{O}_7$

LDA+U method

All-in/All-out magnetic order




$U=0$



$U=1.5$ eV

enlarge



$U=2.0$ eV



Wan et al., (2011)

U



+U calculation

Weyl Semimetal



Wan, Turner, Vishwanath, Savrasov, Phys. Rev. B **83**, 205101 (2011)

Weyl Semimetal

TaAs:

Theory:

Weng et al., PRX (2015)

Huang et al., Nature Commum. (2015)

Exp:


Lv et al., PRX (2015), Nature Phys. (2015);

Xu et al., Science (2015);



.....

Weyl Semimetal Phase in Noncentrosymmetric Transition-Metal Monophosphides

Hongming Weng,^{1,2,*} Chen Fang,³ Zhong Fang,^{1,2} B. Andrei Bernevig,⁴ and Xi Dai^{1,2}




3D view of the nodal rings (in the absence of SOC) and Weyl points (with SOC) in the BZ. Once the SOC is turned on, the nodal rings are gapped and give rise to Weyl points off the mirror planes



Berry curvature distribution


Novel Properties of Quasiparticle

➤ **Klein tunneling** (Katsnelson et al., 2006)




➤ **Fermi Arc**

(Wan, Turner, Vishwanath, Savrasov, (2011))



➤ **Chiral Anomaly**

(Aji (2011), Son & Spivak (2013))



Topological (Crystalline) Insulators

➤ Z₂ topological insulator (*Kane&Mele PRL 2005*)

➤ Mirror Chern Insulator

Ando and Fu, Annu. Rev. Condens. Matter Phys. 6, 361 (2015)

➤ Hourglass Fermions

Wang, Alexandradinata, Cava, Bernevig, Nature 532, 189 (2016).

➤ Higher-order topological insulators

Fang&Fu;Schindler et al., Langbehn et al., Song, Fang&Fang;Benalcazar et al., (2017)

➤ Quantized electric multipole insulators

Benalcazar, Bernevig, Hughes, Science 357, 61 (2017)

➤ Nodal-chain metals

Bzdušek, Wu, Rüegg, Sigrist & Soluyanov, Nature 538, 75 (2016).

➤ Three-fold (or higher) band degeneracies Semimetal

Bradlyn et al., Science (2016)

Semimetal

- Dirac Semimetal
- Weyl Semimetal
- Multi-degeneracy points
- Nodal-line semimetal
- Hopf-line Semimetal
-

How to find Topo Mater

There are so much topological blablabla

Many topological invariants

➤ **Wave function?!!!**

➤ **Need calculate all** topological invariants for one material

Thus find any new topo-mater is a big success

Crystal Symmetry

- Analyzing various crystallographic symmetries

Fu & Kane (2007); parity \rightarrow Z_2 Topological insulator

Fang, Gilbert & Bernevig, (2012); Slager et al., (2012);

Fang & Fu (2015); Fang & Fu (2017); Zhou et al., (2018).....

Do not use wave-function


- Exploiting the mismatch between the real and momentum-space descriptions of the band structure

Kruthoff et al., PRX (2017); Bradlyn et al., Nature (2017);

Po, Vishwanath & Watanabe, Nature Communications (2017)

Topological quantum chemistry

Barry Bradlyn^{1*}, L. Elcoro^{2*}, Jennifer Cano^{1*}, M. G. Vergniory^{3,4,5*}, Zhijun Wang^{6*}, C. Felser⁷, M. I. Aroyo² & B. Andrei Bernevig^{3,6,8,9}



Atomic limit

Representation + Compatibility

Graph theory and band structure


Connected bands (EBR) \rightarrow Semimetal

Disconnected bands (EBR) \rightarrow topological insulator

In total 10403 different EBRs



Topological semimetal



Topological insulator

Symmetry-based Indicators of Band Topology in the 230 Space Groups

Hoi Chun Po,^{1,2} Ashvin Vishwanath,^{1,2,*} and Haruki Watanabe³

Band structure (has direct band gap at high symmetry points)

Atomic limit

$$\{\text{BS}\} \equiv \ker \mathcal{C} \cap \mathbb{Z}^D \simeq \mathbb{Z}^{d_{\text{BS}}} \quad \text{BS} = \sum_{i=1}^{d_{\text{BS}}} m_i \mathbf{b}_i.$$

Quotient Group

Provide no insight into how to find or engineer materials in any non-trivial class

$$\{\text{AI}\} \simeq \mathbb{Z}^{d_{\text{AI}}} \equiv \left\{ \sum_{i=1}^{d_{\text{AI}}} m_i \mathbf{a}_i : m_i \in \mathbb{Z} \right\}$$

商群

$$X_{\text{BS}} \equiv \frac{\{\text{BS}\}}{\{\text{AI}\}}$$

X_{BS}	Space groups
\mathbb{Z}_2	81, 82, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 215, 216, 217, 218, 219, 220
\mathbb{Z}_3	188, 190
\mathbb{Z}_4	52, 56, 58, 60, 61, 62, 70, 88, 126, 130, 133, 135, 136, 137, 138, 141, 142, 163, 165, 167, 202, 203, 205, 222, 223, 227, 228, 230
\mathbb{Z}_8	128, 225, 226
\mathbb{Z}_{12}	176, 192, 193, 194
$\mathbb{Z}_2 \times \mathbb{Z}_4$	14, 15, 48, 50, 53, 54, 55, 57, 59, 63, 64, 66, 68, 71, 72, 73, 74, 84, 85, 86, 125, 129, 131, 132, 134, 147, 148, 162, 164, 166, 200, 201, 204, 206, 224
$\mathbb{Z}_2 \times \mathbb{Z}_8$	87, 124, 139, 140, 229
$\mathbb{Z}_3 \times \mathbb{Z}_3$	174, 187, 189
$\mathbb{Z}_4 \times \mathbb{Z}_8$	127, 221
$\mathbb{Z}_6 \times \mathbb{Z}_{12}$	175, 191
$\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_4$	11, 12, 13, 49, 51, 65, 67, 69
$\mathbb{Z}_2 \times \mathbb{Z}_4 \times \mathbb{Z}_8$	83, 123
$\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_4$	2, 10, 47

Our Algorithm

Space group

symmetry analysis

General atomic basis
 $\{\mathbf{a}_i\}$


Material-specific representations
 \mathbf{n}

linear expansion

$$\mathbf{n} = \sum_i q_i \mathbf{a}_i$$


inspect $\{q_i\}$

Case 1




No topology detected

Case 2



Topological (crystalline) insulator

Case 3



Topological (semi-)metals

$$\mathbf{n} = (\nu, n_{\mathbf{k}_1}^1, \dots, n_{\mathbf{k}_1}^{\alpha_1}, n_{\mathbf{k}_2}^1, \dots, n_{\mathbf{k}_2}^{\alpha_2}, \dots, n_{\mathbf{k}_N}^1, \dots, n_{\mathbf{k}_N}^{\alpha_N})$$

230 space group

Go beyond one-by-one

Case 1: the expansion coefficients q_i 's are all integers, which indicates that \mathbf{n} is an AI, thus the material is equivalent to an atomic insulator;

Case 2: the expansion coefficients q_i 's are not all integers, but all $q_i C_i$'s are integers. This indicates that the \mathbf{n} is a BS and the material is equivalent to a BI, but this system must be topological since some remainder(s) should be nonzero as shown in Eq. 3, i.e. the SI is non-vanishing;

Case 3: Not only the expansion coefficients q_i 's but also the $q_i C_i$'s are not all integers:


(1) some $n_{\mathbf{k}}^\alpha$ is non-integer which indicates that band crossing happens at this k point;

(2) all the $n_{\mathbf{k}}^{\alpha'}$'s are integers, then there must be band crossing in high symmetry line or plane.

*Tang, Po, Vishwanath, Wan**
15, 470 Nature Physics (2019)

Band Structure

**Band structure of Si
(SG227)**



Topological is Global

What is detail:
If nothing pass the Fermi level, then all is detail!

BZ

《物理》 48, 341 (2019)

Bands can cross when they carry different symmetry labels


Dimensions of the irreps determine how the bands are “stuck”

高对称点	坐标	Irreps
Γ	(0, 0, 0)	G_{96}^8 : 4(2), 5(2), 8(4), 12(2), 13(2), 16(4)
X	(0, 1, 0)	G_{64}^2 : 19(4)
L	(1/2, 1/2, 1/2)	G_{24}^3 : 3(1), 4(1), 6(2), 9(1), 10(1), 12(2)
W	(1/2, 1, 0)	G_{64}^3 : 13(1), 14(1), 15(1), 16(1), 20(2)

High symmetry point, High symmetry line, irreps,

Band Structure

Band structure of Si



What is detail:

If nothing pass the Fermi level, then all is detail!

$$\mathbf{n}_{\text{BS}} = \left(\nu, n_{k_1}^1, n_{k_1}^2, \dots, n_{k_1}^{r_1}, n_{k_2}^1, n_{k_2}^2, \dots, n_{k_2}^{r_2}, \dots, n_{k_N}^1, n_{k_N}^2, \dots, n_{k_N}^{r_N} \right).$$

$$\mathbf{n}_{\text{BS}} = (8, 1, 1, 1, 0, 0, 0, 2, 0, 0, 1, 1, 1, 2, 1, 1, 1, 1, 2).$$

$$\mathbf{n}_{\text{BS}, 1} = (4, 0, 0, 1, 0, 0, 0, 1, 0, 0, 0, 1, 1, 1, 0, 0, 1, 1, 1),$$

$$\mathbf{n}_{\text{BS}, 2} = (4, 1, 1, 0, 0, 0, 0, 1, 0, 0, 1, 0, 0, 0, 1, 0, 0, 1, 1, 0, 0, 1).$$


高对称点	坐标	Irreps
Γ	(0, 0, 0)	G_{96}^8 : 4(2), 5(2), 8(4), 12(2), 13(2), 16(4)
X	(0, 1, 0)	G_{64}^2 : 19(4)
L	(1/2, 1/2, 1/2)	G_{24}^3 : 3(1), 4(1), 6(2), 9(1), 10(1), 12(2)
W	(1/2, 1, 0)	G_{64}^3 : 13(1), 14(1), 15(1), 16(1), 20(2)

$$\mathbf{n}_{\text{BS}} = \mathbf{n}_{\text{BS}, 1} + \mathbf{n}_{\text{BS}, 2}$$

➤ Labels become simple counting!

➤ Gap conditions above and below ensure counting is well defined

Atomic limit



- Unit cell has one atom
- One s orbit at this atom


230 Space group

- Wyckoff Positions
- Site symmetry

Wyckoff Positions of Group 2 (P -1)

Multiplicity	Wyckoff letter	Site symmetry	Coordinates
2	i	1	(x, y, z) (-x, -y, -z)
1	h	-1	(1/2, 1/2, 1/2)
1	g	-1	(0, 1/2, 1/2)
1	f	-1	(1/2, 0, 1/2)
1	e	-1	(1/2, 1/2, 0)
1	d	-1	(1/2, 0, 0)
1	c	-1	(0, 1/2, 0)
1	b	-1	(0, 0, 1/2)
1	a	-1	(0, 0, 0)

Wyckoff Positions of Group 227 (F d -3 m) [origin choice



Multiplicity	Wyckoff letter	Site symmetry	Coordinates					
			(0,0,0)+ (0,1/2,1/2)+ (1/2,0,1/2)+ (1/2,1/2,0)+					
192	i	1	(x, y, z)	(-x, -y + 1/2, z + 1/2)	(-x + 1/2, y + 1/2, -z)			
			(z, x, y)	(z + 1/2, -x, -y + 1/2)	(-z, -x + 1/2, y + 1/2)			
			(y, z, x)	(-y + 1/2, z + 1/2, -x)	(y + 1/2, -z, -x + 1/2)			
			(y + 3/4, x + 1/4, -z + 3/4)	(-y + 1/4, -x + 1/4, -z + 1/4)	(y + 1/4, -x + 3/4, z + 3/4)			
			(x + 3/4, z + 1/4, -y + 3/4)	(-x + 3/4, z + 3/4, y + 1/4)	(-x + 1/4, -z + 1/4, -y + 1/4)			
			(z + 3/4, y + 1/4, -x + 3/4)	(z + 1/4, -y + 3/4, x + 3/4)	(-z + 3/4, y + 3/4, x + 1/4)			
			(-x + 1/4, -y + 1/4, -z + 1/4)	(x + 1/4, y + 3/4, -z + 3/4)	(x + 3/4, -y + 3/4, z + 1/4)			
			(-z + 1/4, -x + 1/4, -y + 1/4)	(-z + 3/4, x + 1/4, y + 3/4)	(z + 1/4, x + 3/4, -y + 3/4)			
			(-y + 1/4, -z + 1/4, -x + 1/4)	(y + 3/4, -z + 3/4, x + 1/4)	(-y + 3/4, z + 1/4, x + 3/4)			
			(-y + 1/2, -x, z + 1/2)	(y, x, z)	(-y, x + 1/2, -z + 1/2)			
			(-x + 1/2, -z, y + 1/2)	(x + 1/2, -z + 1/2, -y)	(x, z, y)			
			(-z + 1/2, -y, x + 1/2)	(-z, y + 1/2, -x + 1/2)	(z + 1/2, -y + 1/2, -x)			
96	h	..2	(1/8, y, -y + 1/4)	(7/8, -y + 1/2, -y + 3/4)	(3/8, y + 1/2, y + 3/4)	(5/8, -y, y + 1/4)		
			(-y + 1/4, 1/8, y)	(-y + 3/4, 7/8, -y + 1/2)	(y + 3/4, 3/8, y + 1/2)	(y + 1/4, 5/8, -y)		
			(y, -y + 1/4, 1/8)	(-y + 1/2, -y + 3/4, 7/8)	(y + 1/2, y + 3/4, 3/8)	(-y, y + 1/4, 5/8)		
			(1/8, -y + 1/4, y)	(3/8, y + 3/4, y + 1/2)	(7/8, -y + 3/4, -y + 1/2)	(5/8, y + 1/4, -y)		
			(y, 1/8, -y + 1/4)	(y + 1/2, 3/8, y + 3/4)	(-y + 1/2, 7/8, -y + 3/4)	(-y, 5/8, y + 1/4)		
			(-y + 1/4, y, 1/8)	(y + 3/4, y + 1/2, 3/8)	(-y + 3/4, -y + 1/2, 7/8)	(y + 1/4, -y, 5/8)		
			96	g	..m	(x, x, z)	(-x, -x + 1/2, z + 1/2)	(-x + 1/2, x + 1/2, -z)
						(z, x, x)	(z + 1/2, -x, -x + 1/2)	(-z, -x + 1/2, x + 1/2)
(x, z, x)	(-x + 1/2, z + 1/2, -x)	(x + 1/2, -z, -x + 1/2)						
(x + 3/4, x + 1/4, -z + 3/4)	(-x + 1/4, -x + 1/4, -z + 1/4)	(x + 1/4, -x + 3/4, z + 3/4)						

Pyrochlore structure: Space Group 227

Y → 16d

Ir → 16c

O1 → 8b

O2 → 48f

Calculation of Atomic Insulator Basis

Taking SG2 as an example

Step 1:

Obtaining HSPs.

$SG2's$ HSP	Γ	X	Y	Z	U	T	S	R
coordinate	(0,0,0)	($\frac{1}{2}$, 0, 0)	(0, $\frac{1}{2}$, 0)	(0, 0, $\frac{1}{2}$)	($\frac{1}{2}$, $\frac{1}{2}$, 0)	(0, $\frac{1}{2}$, $\frac{1}{2}$)	($\frac{1}{2}$, 0, $\frac{1}{2}$)	($\frac{1}{2}$, $\frac{1}{2}$, $\frac{1}{2}$)

Step 2: Give all the Wyckoff positions.

2  8  1  17

$SG2's$ Wyckoff position	site-group	Wyckoff orbits
2i	C_1	$(x, y, z), (-x, -y, -z)$
1h	C_i	$(1/2, 1/2, 1/2)$
1g	C_i	$(0, 1/2, 1/2)$
1f	C_i	$(1/2, 0, 1/2)$
1e	C_i	$(1/2, 1/2, 0)$
1d	C_i	$(1/2, 0, 0)$
1c	C_i	$(0, 1/2, 0)$
1b	C_i	$(0, 0, 1/2)$
1a	C_i	$(0, 0, 0)$

TABLE VII. The nine Wyckoff positions for $SG2$.

Calculation of Atomic Insulator Basis

$SG2$	n_{2i}^1	n_{1h}^1	n_{1h}^2	n_{1g}^1	n_{1g}^2	n_{1f}^1	n_{1f}^2	n_{1e}^1	n_{1e}^2	n_{1d}^1	n_{1d}^2	n_{1c}^1	n_{1c}^2	n_{1b}^1	n_{1b}^2	n_{1a}^1	n_{1a}^2
ν	4	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2
Γ_1^1	1	1	0	1	0	1	0	1	0	1	0	1	0	1	0	1	0
Γ_2^1	1	0	1	0	1	0	1	0	1	0	1	0	1	0	1	0	1
X_1^1	1	1	0	0	1	1	0	0	1	0	1	0	1	1	0	1	0
X_2^1	1	0	1	1	0	0	1	1	0	1	0	1	0	0	1	0	1
Y_1^1	1	1	0	0	1	0	1	1	0	0	1	1	0	0	1	1	0
Y_2^1	1	0	1	1	0	1	0	0	1	1	0	0	1	1	0	0	1
Z_1^1	1	1	0	0	1	0	1	0	1	1	0	1	0	1	0	0	1
Z_2^1	1	0	1	1	0	1	0	1	0	0	1	0	1	0	1	1	0
U_1^1	1	1	0	1	0	0	1	0	1	1	0	0	1	0	1	1	0
U_2^1	1	0	1	0	1	1	0	1	0	0	1	1	0	1	0	0	1
T_1^1	1	1	0	1	0	1	0	0	1	0	1	1	0	0	1	0	1
T_2^1	1	0	1	0	1	0	1	1	0	1	0	0	1	1	0	1	0
S_1^1	1	1	0	1	0	0	1	1	0	0	1	0	1	1	0	0	1
S_2^1	1	0	1	0	1	1	0	0	1	1	0	1	0	0	1	1	0
R_1^1	1	1	0	0	1	1	0	1	0	1	0	0	1	0	1	0	1
R_2^1	1	0	1	1	0	0	1	0	1	0	1	1	0	1	0	1	0

TABLE X. The 17 AI vectors for $SG2$.

$n = \{ \nu, n_{\alpha_1}^1, n_{\alpha_2}^2, n_X^1, n_X^2, n_Y^1, n_Y^2, n_Z^1, n_Z^2, n_U^1, n_U^2, n_T^1, n_T^2, n_S^1, n_S^2, n_R^1, n_R^2 \}$

The BS can be represented by an integer-valued vector, $\mathbf{n} = (\nu, n_{\mathbf{k}_1}^1, n_{\mathbf{k}_1}^2, \dots, n_{\mathbf{k}_1}^{\alpha_1}, \dots, n_{\mathbf{k}_1}^{r_1}, n_{\mathbf{k}_2}^1, n_{\mathbf{k}_2}^2, \dots, n_{\mathbf{k}_2}^{\alpha_2}, \dots, n_{\mathbf{k}_2}^{r_2},$

$n_{\mathbf{k}_i}^1, \dots, n_{\mathbf{k}_i}^{\alpha_i}, \dots, n_{\mathbf{k}_i}^{r_i}, \dots, n_{\mathbf{k}_N}^1, \dots, n_{\mathbf{k}_N}^{\alpha_N}, \dots, n_{\mathbf{k}_N}^{r_N})$.

Calculation of Atomic Insulator Basis

We choose our basis such that the values of C_i are maximized: **Smith decomposition**.

(a) The 9 AI basis vectors for $SG2$. Here ν is the number of the bands. It is also called the filling number. Starting from the 3rd row, we give the number $n_{\mathbf{k}}^\alpha$ in order. We omit the notation n for clarity: The first column of these rows gives the information of the HSP and its irrep completely.

$$Z_2 \quad \triangleleft Z_2 \quad \triangleleft Z_2 \quad \triangleleft Z_4$$

$$BS = \sum_{i=1}^{d_{BS}} m_i \mathbf{b}_i,$$

Common factors

$$d_{AI} = d_{BS} \quad \text{Po et al., NC (2017)}$$

$SG2$	a_1	a_2	a_3	a_4	a_5	a_6	a_7	a_8	a_9
ν	4	2	2	2	2	4	4	4	8
Γ_1^1	1	1	1	1	1	2	2	2	4
Γ_2^1	1	0	0	0	0	0	0	0	0
X_1^1	1	1	0	1	0	2	2	0	4
X_2^1	1	0	1	0	1	0	0	2	0
Y_1^1	1	1	0	0	1	2	0	2	4
Y_2^1	1	0	1	1	0	0	2	0	0
Z_1^1	1	1	0	0	0	0	0	0	0
Z_2^1	1	0	1	1	1	2	2	2	4
U_1^1	1	1	1	0	0	0	2	2	4
U_2^1	1	0	0	1	1	2	0	0	0
T_1^1	1	1	1	1	0	2	2	2	4
T_2^1	1	0	0	0	1	0	0	0	0
S_1^1	1	1	1	0	1	2	2	2	4
S_2^1	1	0	0	1	0	0	0	0	0
R_1^1	1	1	0	1	1	2	2	2	4
R_2^1	1	0	1	0	0	0	0	0	0

Our Algorithm

Space group

symmetry analysis

General atomic basis
 $\{\mathbf{a}_i\}$


Material-specific representations
 \mathbf{n}

linear expansion

$$\mathbf{n} = \sum_i q_i \mathbf{a}_i$$


inspect $\{q_i\}$

Case 1




No topology detected

Case 2



Topological (crystalline) insulator

Case 3



Topological (semi-)metals

$$\mathbf{n} = (\nu, n_{\mathbf{k}_1}^1, \dots, n_{\mathbf{k}_1}^{\alpha_1}, n_{\mathbf{k}_2}^1, \dots, n_{\mathbf{k}_2}^{\alpha_2}, \dots, n_{\mathbf{k}_N}^1, \dots, n_{\mathbf{k}_N}^{\alpha_N})$$

230 space group
Go beyond one-by-one

Case 1: the expansion coefficients q_i 's are all integers, which indicates that \mathbf{n} is an AI, thus the material is equivalent to an atomic insulator;

Case 2: the expansion coefficients q_i 's are not all integers, but all $q_i C_i$'s are integers. This indicates that the \mathbf{n} is a BS and the material is equivalent to a BI, but this system must be topological since some remainder(s) should be nonzero as shown in Eq. 3, i.e. the SI is non-vanishing;


Case 3: Not only the expansion coefficients q_i 's but also the $q_i C_i$'s are not all integers:

(1) some $n_{\mathbf{k}}^\alpha$ is non-integer which indicates that band crossing happens at this k point;

(2) all the $n_{\mathbf{k}}^\alpha$'s are integers, then there must be band crossing in high symmetry line or plane.

*Tang, Po, Vishwanath, Wan**
15, 470 Nature Physics (2019)

\mathbb{Z}_4 TCI



2 of \mathbb{Z}_4

TABLE I. We focus on the following space groups (\mathcal{SG} s), in which a strong topological insulators generate a \mathbb{Z}_4 subgroup in the group of symmetry indicators, X_{BS} . The entry $2 \in \mathbb{Z}_4$ corresponds to various kinds of topological crystalline insulators, and the predicted materials candidates for such phases are tabulated.

X_{BS}	$\mathbb{Z}_2^3 \times \mathbb{Z}_4$	$\mathbb{Z}_2^2 \times \mathbb{Z}_4$	$\mathbb{Z}_2 \times \mathbb{Z}_4$	\mathbb{Z}_4
\mathcal{SG}	2	11, 12	166	61, 136, 227
Materials	Ag_2F_5	$\beta\text{-MoTe}_2, \text{BiBr}$	A7-P	c-TiS ₂

Topological Hinge states in β -MoTe₂

(SG) 11 ($P2_1/m$).

α -phase:

β -phase: room temperature

Monolayer—QSH (*Qian, Liu, Fu & Li, Science 2014*)

γ -phase:



Type-II Weyl Semimetal:

Deng et al., Nature Phys. (2016)

Tamai et al., PRX (2016)

Huang et al., Nature Mater. (2016)

Jiang et al., Nature Commun. (2016)



DFT results of β -MoTe₂

- 2 inequivalent Mo's and 4 in-equivalent Te's
 - all occupy 2e Wyckoff positions.
 - total 12 atoms (56 valence electrons) primitive unit cell
- v=56** **SG11**

TABLE I. For $SG11$, the HSPs are given by the labels Γ, B, \dots in order. For the labeling of the irreps of $\mathcal{G}(\mathbf{k}_i)$, we use (j, m) where j means the j th irrep and m denotes the dimension of the corresponding irrep. They are all listed in Ref. [2]. We use the same order of the irrep as Ref. [2]. The red color means that due to \mathcal{T} , the irrep must occur with its \mathcal{T} pair (belonging to the same irrep) simultaneously. Thus \mathcal{T} requires that the red colored irreps must happen even times. So it is necessary to divide them by 2 [1] to obtain the physical common factors.

HSP	Γ				B				Y				Z	C	D	A				E
irrep	(1,1)	(2,1)	(3,1)	(4,1)	(1,1)	(2,1)	(3,1)	(4,1)	(1,1)	(2,1)	(3,1)	(4,1)	(1,2)	(1,2)	(1,2)	(1,1)	(2,1)	(3,1)	(4,1)	(1,2)
$n_{\mathbf{k}_i}^{\alpha_i}$	16	16	12	12	14	14	14	14	14	14	14	14	14	14	14	14	14	14	14	14

$$\mathbf{n} = (\nu, n_{\mathbf{k}_1}^1, n_{\mathbf{k}_1}^2, \dots, n_{\mathbf{k}_1}^{\alpha_1}, \dots, n_{\mathbf{k}_2}^1, n_{\mathbf{k}_2}^2, \dots, n_{\mathbf{k}_2}^{\alpha_2}, \dots, n_{\mathbf{k}_N}^1, n_{\mathbf{k}_N}^2, \dots, n_{\mathbf{k}_N}^{\alpha_N}, \dots),$$

DFT



$$\mathbf{n} = (56, n_{\mathbf{k}_1}^1, n_{\mathbf{k}_1}^2, n_{\mathbf{k}_1}^3, n_{\mathbf{k}_1}^4, n_B^1, n_B^2, n_B^3, n_B^4, n_Y^1, n_Y^2, n_Y^3, n_Y^4, n_Z^1, n_C^1, n_D^1, n_A^1, n_A^2, n_A^3, n_A^4, n_E^1)$$

$$\mathbf{n} = (56, 16, 16, 12, 12, 14, 14, 14, 14, 14, 14, 14, 14, 14, 14, 14, 14, 14, 14, 14, 14)$$

Expansion of β -MoTe₂

(b) The 5 AI basis vectors for $SG11$.

$SG11$	\mathbf{a}_1	\mathbf{a}_2	\mathbf{a}_3	\mathbf{a}_4	\mathbf{a}_5
ν	4	4	0	0	0
Γ_1^1	1	2	0	0	0
Γ_2^1	1	2	0	0	0
Γ_3^1	1	0	0	0	0
Γ_4^1	1	0	0	0	0
B_1^1	1	0	0	2	0
B_2^1	1	0	0	2	0
B_3^1	1	2	0	-2	0
B_4^1	1	2	0	-2	0
Y_1^1	1	0	2	0	0
Y_2^1	1	0	2	0	0
Y_3^1	1	2	-2	0	0
Y_4^2	1	2	-2	0	0
Z_1^2	1	1	0	0	0
C_1^2	1	1	0	0	0
D_1^2	1	1	0	0	0
A_1^1	1	2	-2	-2	4
A_2^1	1	2	-2	-2	4
A_3^1	1	0	2	2	-4
A_4^1	1	0	2	2	-4
E_1^2	1	1	0	0	0

$$C_1 = C_2 = 1, C_3 = C_4 = 2, \text{ and } C_5 = 4$$

$$\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_4$$

$$\mathbf{n} = (56, 16, 16, 12, 12, 14, 14, 14, 14, 14, 14, 14, 14, 14, 14, 14, 14, 14, 14, 14)$$

$$= 12\mathbf{a}_1 + 2\mathbf{a}_2 + \mathbf{a}_3 + \mathbf{a}_4 + \frac{1}{2}\mathbf{a}_5,$$

$$(q_1, q_2, q_3, q_4, q_5) = (12, 2, 1, 1, \frac{1}{2})$$

$$SI (0, 0, 2)$$

$$2 \in \mathbb{Z}_4$$

2 of \mathbb{Z}_4

Topological feature of β -MoTe₂

$$\kappa_1 = \sum_{\mathbf{k} \in \text{TRIM}} (n_{\mathbf{k}}^+ - n_{\mathbf{k}}^-) / 2 \pmod{4}$$

Z. Song, T. Zhang, Z. Fang, and C. Fang, ArXiv e-prints (2017), [arXiv:1711.11049](#).
E. Khalaf, H. C. Po, A. Vishwanath, and H. Watanabe, ArXiv e-prints (2017), [arXiv:1711.11589](#).

$\mathbf{k} \in \text{TRIM}$	Γ	X	Y	Z	U	T	S	R
$n_{\mathbf{k}}^+$	16	14	14	14	14	14	14	14
$n_{\mathbf{k}}^-$	12	14	14	14	14	14	14	14

$n_{\mathbf{k}}^{\pm}$ is the number of the occupied even/odd Kramers pairs


For this compound, 2 of $Z_4 \rightarrow$ mirror Chern/hinge states

WIEN2K \rightarrow mirror Chern number for $k_z = 0$ and $k_z = \pi/c$


They all vanish

So hinge state!


Hinge state from TB calculation




c



TB reproduces topo-feature and band dispersion



d



*Had been confirmed by
Wang, Wieder, Li, Yan & Bernevig,
arXiv:1806.11116 (2018)*

BiBr: Dirac Surface States Coexisting with Hinge State

$SG12$ ($C2/m$)

TABLE III. For $SG12$, the HSPs are given by the labels Γ, A, \dots in order, and their coordinates can be referred to Ref. [2]. For the labeling of the irreps of $\mathcal{G}(\mathbf{k}_i)$, we use (j, m) where j means the j th irrep as listed in order by Ref. [2] and m denotes the dimension. The red color means that due to \mathcal{T} , the irrep must occur simultaneously with its \mathcal{T} pair which belongs to the same irrep.

HSP	Γ				A				Z				M				L		V	
irrep	(1,1)	(2,1)	(3,1)	(4,1)	(1,1)	(2,1)	(3,1)	(4,1)	(1,1)	(2,1)	(3,1)	(4,1)	(1,1)	(2,1)	(3,1)	(4,1)	(1,1)	(2,1)	(1,1)	(2,1)
$n_{\mathbf{k}_i}^{\alpha_i}$	18	18	14	14	16	16	16	16	16	16	16	16	16	16	16	16	16	16	16	16

$$\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_4$$


$$\mathbf{n} = (64, 18, 18, 14, 14, 16, 16, 16, 16, 16, 16, 16, 16, 16, 16, 16, 16, 16, 16, 16)$$

$$= 14\mathbf{a}_1 + 2\mathbf{a}_2 + 2\mathbf{a}_3 + 2\mathbf{a}_4 + \mathbf{a}_6 - \frac{1}{2}\mathbf{a}_7,$$





SI of (0, 0, 2)

(c) The 7 AI basis vectors for $SG12$.

$SG12$	\mathbf{a}_1	\mathbf{a}_2	\mathbf{a}_3	\mathbf{a}_4	\mathbf{a}_5	\mathbf{a}_6	\mathbf{a}_7
ν	4	4	-8	2	-4	8	-8
Γ_1^1	1	2	-4	1	-2	4	-4
Γ_2^1	1	2	-4	1	-2	4	-4
Γ_3^1	1	0	0	0	0	0	0
Γ_4^1	1	0	0	0	0	0	0
A_1^1	1	0	-1	0	0	2	-4
A_2^1	1	0	-1	0	0	2	-4
A_3^1	1	2	-3	1	-2	2	0
A_4^1	1	2	-3	1	-2	2	0
Z_1^1	1	0	-2	1	0	2	-4
Z_2^1	1	0	-2	1	0	2	-4
Z_3^1	1	2	-2	0	-2	2	0
Z_4^1	1	2	-2	0	-2	2	0
M_1^1	1	2	-3	0	-2	4	0
M_2^1	1	2	-3	0	-2	4	0
M_3^1	1	0	-1	1	0	0	-4
M_4^1	1	0	-1	1	0	0	-4
L_1^1	1	1	-1	1	-2	0	0
L_2^1	1	1	-3	0	0	4	-4
V_1^1	1	1	0	0	0	0	0
V_2^1	1	1	-4	1	-2	4	-4



Topological states in BiBr



- Mirror Chern number = 0
- Hinge-State + Surface Dirac State
- TB \rightarrow topo-feature

Confirmed by:


Hsu C H *et al.* 2D Mater., 2019, 6:031004

Non-Centrosymmetric Strong TI in AgNaO

- Inversion \rightarrow Fu-Kane parity criterion
- Non-inversion \rightarrow calculate Z_2 invariant

SG 216 ($F\bar{4}3m$), which are non-centrosymmetric and contains S_4

Expansion \rightarrow be insulators with the SI $1 \begin{matrix} \square \\ \square \end{matrix} Z_2$



$s(d)$ band is mainly above (below) the Fermi level.

band inversion near the Gamma point,

results in an S_4 invariant




Other $2\in Z_4$ Materials

- Cubic crystal TiS_2 (SG227) is a glide-protected TCI with **hourglass** surface states
- Elemental phosphorus in the A7 structure (SG166), which occurs at about 9GPa, is predicted to be an inversion-protected TCI with 1D **hinge** states
- Ag_2F_5 (SG2) is a **weak TI** with additional inversion-protected band topology characterized by the invariant $\kappa_1=2$

Strong TI Found by 1,3 of \mathbb{Z}_4

TABLE V. Table of centrosymmetric STI candidates discovered by 1, 3 in \mathbb{Z}_4

$S\mathcal{G}$	Material	X_{BS}	SI	$(\nu_0; \nu_1, \nu_2, \nu_3)$	κ_1
2	CaAs ₃ [17]	$\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_4$	(0,0,1,1)	(1;1,0,0)	1
166	Bi ₂ PbTe ₄ [18]	$\mathbb{Z}_2 \times \mathbb{Z}_4$	(1,1)	(1;1,1,1)	3
166	CaGa ₂ As ₂ [19]	$\mathbb{Z}_2 \times \mathbb{Z}_4$	(1,1)	(1;1,1,1)	1




Dirac Semimetal MgBi_2O_6

$SG136$


Expansion \rightarrow case 3

So topological Semimetal




e) The 8 AI basis vectors for $SG136$.

$SG136$	a_1	a_2	a_3	a_4	a_5
ν	8	8	8	4	-48
Γ_1^2	1	2	2	1	-12
Γ_2^2	1	0	2	1	-8
Γ_3^2	1	0	0	0	0
Γ_4^2	1	2	0	0	-4
K_1^2	1	1	0	1	-4
K_2^2	1	1	0	1	-4
K_3^2	1	1	2	0	-8
K_4^2	1	1	2	0	-8
K_1^4	2	2	2	1	-12
K_2^4	2	2	2	1	-12
K_1^1	1	1	1	0	-4
K_2^1	1	1	1	0	-4
K_3^1	1	1	1	0	-4
K_4^1	1	1	1	0	-4
K_5^1	1	1	1	1	-8
K_6^1	1	1	1	1	-8
K_7^1	1	1	1	1	-8
K_8^1	1	1	1	1	-8
K_1^2	2	2	2	1	-12
K_2^2	2	2	2	1	-12



Three-Fold Degenerate Fermions

- AuLiMgSn (SG216)
- DFT $\rightarrow n_{\mathbf{k}}^{\alpha}$ are all integers thus there are finite direct gaps in all the HSPs.
- However expansion on the SG216's AI basis vectors shows that they cannot constitute a BS at all, namely case 3 in the main text



The Nodal-line Semimetal AgF_2

AgF_2 (SG61)

$$\left(\frac{11}{2}, \frac{19}{2}, \frac{13}{4} \right)$$

(d) The 3 AI basis vectors for SG61.

SG61	\mathbf{a}_1	\mathbf{a}_2	\mathbf{a}_3
ν	16	8	-32
Γ_1^2	4	4	-12
Γ_2^2	4	0	-4
Y_1^2	4	2	-8
Y_2^2	4	2	-8
X_1^2	4	2	-8
X_2^2	4	2	-8
Z_1^2	4	2	-8
Z_2^2	4	2	-8
U_1^2	2	1	-4
U_2^2	2	1	-4
T_1^2	2	1	-4
T_2^2	2	1	-4
S_1^2	2	1	-4
S_2^2	2	1	-4
R_1^1	1	0	0
R_2^1	1	0	0
R_3^1	1	0	0
R_4^1	1	0	0
R_5^1	1	1	-4
R_6^1	1	1	-4
R_7^1	1	1	-4
R_8^1	1	1	-4


Comprehensive database searches

- Around 180000 compounds at ICSD database
- Consider stoichiometric compound
- Do not consider $3d/4f/5f$ and several $4d/5d$ element
- ➔ **19143** compounds with SOC+time-reverse symmetry

- Highly efficient (main jobs is around 1 month within a cluster 28-node/one-node-32 CPU core)

GGA calculation

Check by MBJ calculation



List 258 good topological insulator

ccmp.nju.edu.cn

$S\bar{G}$	X_{BS}	Topological insulators
1	$Z_2^3 \times Z_4$	AgO ₄ S[3], Ag₂Fe [4], As ₂ Ca[5, 6], As ₂ Sr[7, 8], Bi[9], Br ₅ W[10], CaMo ₆ S ₈ [11], CaP ₂ [8, 12]
2	$Z_2^2 \times Z_4$	BBeLi[13], GeHPd[14], HfS ₃ [15], Mo ₂ S ₃ [16], NbPt ₃ [17], Nb ₂ Se ₃ [18], Se ₂ Ta ₂ [18, 19]
12	$Z_2^2 \times Z_4$	Ag ₂ K ₂ Se ₂ [20], Al ₂ Ge ₄ Sr ₃ [21], Al ₃ Mo ₃ [22], As ₂ Ba[8, 23], As ₂ Sr[8, 24], Au ₂ P ₃ [25], BaSb ₃ [26] Ba ₂ Cd ₂ Sb ₂ [27], Ba ₂ Hg ₂ O ₁₄ Pd[28], Ba ₃ Li ₄ Sn ₃ [29], BaTl₄ [30], Ba ₁₁ Bi ₁₄ Cd ₅ [31], Bi ₂ Pb ₂ Se ₆ [32], Bi ₂ Pd[33] Bi ₄ Pb ₇ Se ₁₃ [34], Br ₄ Cs ₂ I ₂ Pd[35], Hg₂Sn₂Sr [36], Nb ₂ PdS ₅ [37], Nb ₂ PdS ₆ [38], Nb₂PdSe₆ [38], P ₃ Sr[39] , PdSe ₂ Ta ₂ [38], PtSe ₂ Ta ₂ [40]
14	$Z_2 \times Z_4$	Ag ₂ Te[41, 42]
51	$Z_2^2 \times Z_4$	AlPt ₂ [43], AuTe ₃ [44], Au ₂ Rb ₂ Tl[45]
55	$Z_2 \times Z_4$	Al₂Bi₆Ca₈ [46], Bi₆In₂Sr₅ [47]
57	$Z_2^2 \times Z_4$	STa ₂ [48]
58	Z_4	Bi ₂ Hf[49], S ₂ Ti[50]
59	$Z_2 \times Z_4$	Ag ₃ Sb[51], AuCs ₃ Pb ₄ [52], AuPb ₄ Rb ₃ [52], AuRb ₃ Sn ₄ [52]
60	Z_4	Au ₂ Pb[53, 54], O ₂ Pb[55]
61	Z_4	AuSn ₂ [56], Bi ₃ Pt[57]
62	Z_4	AsCdNa[58], As ₂ Hf[59], As ₂ Zr[60, 61], Ba ₂ Pb[62], Ba ₂ Si[63, 64], Ca ₃ GeO[65], CdGeSr[66], CdNaSb[67, 68], FSeY[69] GaPtSc[70], GaPtY[70], GeHPt[71, 72], GeMoZr[73], GePdZr[74, 75], GePtZr[75, 76], GeZr[77], Ge ₂ InLiSr ₂ [78], Ge ₂ Mo[79] Ge ₃ W[80], HfNbP[81], HfP ₂ [82], HfSi[83], NNa ₃ [84], N ₂ Ta ₃ [85], NbPZr[86], PPTc[75, 87], P ₂ Ti[88] P ₂ Zr[89], PbSr ₂ [90, 91], PdSrTi[75, 92], PdSiZr[93], PtSiTi[75, 94], PtSiZr[75, 94], Sb ₂ SrZn[95, 96], SiSr[97], SiTi[98], SiZr[98]
63	$Z_2 \times Z_4$	Ag ₂ Ta[99], AlBMo[100], AlBW[100], Al ₂ BaSn ₂ [101], AuClTe[102], B ₇ WY ₃ [103], BaIn[104], CaSi [105] Ga[106], GeNa ₂ Zn[107], HfPd[108], HfTe ₅ [109, 110], PdZr[111]
64	$Z_2 \times Z_4$	As[112], C ₂ B ₂ Mg[113], LiTl[114]
65	$Z_2^2 \times Z_4$	Ag₃Te₂Tl [115], Ba ₃ Ge ₄ [116]
69	$Z_2^2 \times Z_4$	B ₂ Zn[117]
71	$Z_2 \times Z_4$	AsTeTi[118], Br ₂ Ca ₃ Si[119]
72	$Z_2 \times Z_4$	Br ₂ Hg ₅ O ₄ [120]
87	$Z_2 \times Z_8$	As ₄ Mo ₅ [121], Hf ₅ Te ₄ [122], Se ₄ Ti ₅ [123], Te ₄ Zr ₅ [124]
114	Z_2	Pd ₄ S[125]
121	Z_2	Ag ₂ S ₄ SnZn[126]
122	Z_2	As ₂ CdGe[127], As ₂ CdSn[128], As ₂ SnZn[129], CdSb ₂ Sn[130]
123	$Z_2 \times Z_4 \times Z_8$	BaGe ₂ Mg ₄ [131], BiLi [132], BiNa[133], ClNa ₂ [134], ClNa ₃ [134]
127	$Z_4 \times Z_8$	B ₄ Y[135], CsI ₃ Sn[136, 137], Pt ₃ Si[138]
129	$Z_2 \times Z_4$	HfSb ₂ [139]
136	Z_4	Ag ₂ CsSe ₃ [140], Ag ₅ CsTe ₃ [141]
139	$Z_2 \times Z_8$	Ag ₂ Zr[142], Au ₂ Hf[143], Au ₂ Zr[144], BaCd ₂ Ge ₂ [145], BaGe ₂ Mg ₂ [146], Be ₁₂ W [147] Bi ₁₀ Ca ₁₁ [148], CaGe ₂ Zn ₂ [146], Cd ₂ Ge ₂ Sr[149], Ge ₂ SrZn ₂ [150], Hf ₂ Pd[151], InPd ₃ [152], PdZr ₂ [153], Pd ₂ Ti[154]
140	$Z_2 \times Z_8$	Bi ₂ In ₅ [155], Hf ₂ Si[156], In ₂ Sb ₂ [157], Pb ₂ Pt[158], Sc[159], SiZr ₂ [160], Sb₂Tl [161], Si ₂ Sr ₅ [162], Sn ₂ Sr ₅ [163]
164	$Z_2 \times Z_4$	Ag ₂ O[164], As ₂ Ge ₂ Te ₃ [165], As ₄ GeTe ₇ [166], BaSn ₂ [167, 168], Be[169], Bi ₂ Li ₃ Y[170], Bi ₂ Mg ₂ [171, 172] Bi ₂ Pb ₂ Se ₆ [173, 174], Bi ₄ SnTe ₇ [175], CNb ₂ [176], CTa ₂ [177], C ₆ K ₂ N ₆ PtS ₆ [178], C ₆ N ₆ PtRb ₂ Se ₆ [178] CaGe ₂ [179, 180], CaSi ₂ [180, 181], GeSb ₄ Te ₇ [182], Ge ₂ Sr[180, 183], PdTe ₂ [184, 185], PtTe ₂ [186, 187], Sb ₂ Te ₃ [188], SiTe ₂ [189]
166	$Z_2 \times Z_4$	CsGa ₇ [190], Cl ₃ Na ₂ Ti ₃ [191], Ga₇Rb [192], PtTe[193], Pt ₂ Te ₃ [193], STi[194], SeTi[195], AgGeLi ₂ [196] Al ₁₀ Ba ₇ [197], As[198], As ₂ CaGa ₂ [199] As ₂ Ge ₄ Te ₇ [200], As ₂ Sn ₂ Sr[201, 202], Bi ₂ GeTe ₄ [203, 204], Bi ₂ PbTe ₄ [205], Bi ₂ SeTe ₂ [206, 207], Bi ₂ Se ₃ [208, 209], Bi ₂ Te ₃ [209, 210] Bi ₆ PbTe ₁₀ [210], Bi ₈ Te ₉ [211], CS₂Ta₂ [212], CY ₂ [213], C ₅ Ti ₈ [214] CaGe ₂ [215], CaSi ₂ [216, 217], GaGeTe[218] GaP[219], Ga ₂ Te ₃ [220], In ₂ Te ₃ [220], Li ₃ Pb ₃ [221] N ₂ W[222], Sb[223], SbTe ₂ Tl[224, 225], Sb ₂ SeTe ₂ [226, 227] Sb ₂ SnTe ₄ [228, 229], Sb ₂ Te ₃ [209, 230], Sb ₅ Te ₃ [231], Sb ₄ Te ₉ [231]
191	$Z_6 \times Z_{12}$	B ₂ Sr[232], BaSi ₂ [233], CaSi ₂ [234], Li ₃ Pd[235], Li ₂ Pt[236], Ti[237]
193	Z_{12}	Pb ₂ TeZr ₅ [238], Sn ₃ Zr ₅ [239]
194	Z_{12}	AgSbSr[240], AlN ₃ Ti ₄ [241], Al ₂ Hf[242], AuBaBi[243], C ₂ AlTa ₃ [244], Ca ₂ IN[245] MgPo[246], PbPt[247], SiSr ₂ [248], BeHfSi[249]
216	Z_2	AgKO[250], AgNaO[250], AgORb[250], HgS[251], InN[252], NITl [253]
221	$Z_4 \times Z_8$	AlSc[254], AlY[255], B ₆ Ca[256], B ₆ Sr[256], BeTi[257] Br ₃ CsGe[258], Br ₃ CsPb[259], CsI ₃ Sn[260] MgPd[261], PbSe[262], SnTe[263]
223	Z_4	BiNb ₃ [264], Nb ₂ Sb[265], SbTa ₃ [266, 267]
225	Z_8	BiY[268], CPd[269], CZr[270], Li ₃ MgSi[271], N ₂ Pd[272]
227	Z_4	CTi ₂ [273], Se[274]

TABLE I. The list of all the TIs with relatively clean Fermi surfaces discovered by GGA calculations: when the MBJ calculation doesn't change the band topology predicted by GGA calculations, we print the material in blue, while the red color means that the MBJ calculation results in a topological crystalline phase.

SG	X_{BS}	Topological crystalline insulators
2	$Z_3^2 \times Z_4$	CsHg[275],HgK[276]
11	$Z_2^2 \times Z_4$	BaSb ₂ [277],MoTe ₂ [278],Sb ₂ Sr[279],Se ₃ Ta[280, 281]
12	$Z_2^3 \times Z_4$	Al ₄ Cl ₅ Zr ₁₂ [282],Al ₄ Na ₄ P ₁₂ Sr ₈ [283],As ₂ Nb[284, 285],As ₂ Ta[285, 286],As ₃ Mo ₂ [287],As ₃ W ₂ [288] As ₄ Ba ₃ Zn ₂ [289],Ba ₃ Cd ₂ Sb ₄ [290],BiBr[291],BiHf ₂ [292],Bi ₂ ITe[293, 294],CdK ₆ Pb ₈ [295],ClZr [296] Cl ₈ NSc ₅ [297],NbP ₂ [298],NbSb ₂ [299],Sb ₂ Ta[300, 301],Se ₄ Ti ₃ [302],Ta ₂ Te ₃ [303]
51	$Z_2^3 \times Z_4$	GaPt ₂ [304],CdMg[305]
55	$Z_2 \times Z_4$	Ca ₅ Ga ₂ Sb ₆ [306]
58	Z_4	Bi ₃ RbS ₅ [307]
59	$Z_2 \times Z_4$	BrNTi[308]
62	Z_4	HgSr ₃ [309],PdSi[310],PdSn[311]
63	$Z_2 \times Z_4$	BaGe[312],BaSi[313], BaSn[314],Ba ₃ Pb ₅ [315],Ba ₅ Cd ₂ FSb ₅ [316], CaGe[317],CaSn [318] Cd ₂ FSb ₅ Sr ₅ [319],PbSr[320],SnSr[321]
64	$Z_2 \times Z_4$	Ca ₅ Ga ₂ N ₄ [322],Li[323]
65	$Z_2^3 \times Z_4$	Au ₇ Rb ₃ [324]
69	$Z_2^3 \times Z_4$	Ge ₆ Li ₂ Sr ₄ [325]
71	$Z_2 \times Z_4$	Ba ₃ Bi ₄ Li ₄ [326],Ba ₃ Li ₄ Sb ₄ [326]
87	$Z_2 \times Z_8$	Au ₄ Ti[327]
88	Z_4	O ₄ PbPd ₂ [328]
123	$Z_2 \times Z_4 \times Z_8$	As ₃ CsZn ₄ [329],As ₃ RbZn ₄ [329],CPd ₃ Sn[330], Pd ₃ Sn[331]
127	$Z_4 \times Z_8$	B ₄ Ca[332]
129	$Z_2 \times Z_4$	AsGeNb[333],GeHfS[334, 335],GeHfSe[334, 335],GeHfTe[334, 335],GeNbSb[336],GeSZr [337] GeSeZr[334],HfSSi[334, 335],HfSeSi[334, 335],OSiZr[334],SSiZr[338],SeSiZr[338],SiTeZr[338],SnTeZr[338]
137	Z_4	Ba ₂ LiN[339]
139	$Z_2 \times Z_8$	AsBa ₂ [340],AsCa ₂ [341] Ba ₂ Bi[342],Ba ₂ Sb[343],Ba ₁₁ Bi ₁₀ [344],BiSr ₂ [343],Bi ₂ F ₂ OSr ₂ Ti ₂ [345],Br ₂ Ca ₃ Si [346] CaIn[347],F ₂ OSb ₂ Sr ₂ Ti ₂ [348],Hf ₂ Hg[349],Pd ₂ Si ₂ Sr[350],SbSr ₂ [351]
140	$Z_2 \times Z_8$	Bi[352],GePt ₃ [353],Mg ₂ Pt[354]
164	$Z_2 \times Z_4$	BaSi ₂ [355],BiTe[356],Bi ₂ Pb ₂ Te ₅ [357, 358], Bi ₂ Se ₂ [359],Bi ₃ STe ₂ [360],HgPt ₂ Se ₃ [361],Te ₂ Zr[362]
166	$Z_2 \times Z_4$	B ₂ Mo[363],Ba ₃ O ₅ Sc ₄ [364],Pt ₃ Te ₄ [365],TeZr ₃ [366],AsNaTe ₂ Zr ₂ [367],As ₃ Cd ₄ K[368] As ₃ Cd ₄ Na[368],As ₃ Cd ₄ Rb[368],Bi[369],Bi ₄ Se ₃ [370, 371],Bi ₈ Se ₉ [372],BrZr[373],CNb ₂ S ₂ [374],C ₃ Al ₆ N ₂ [375] ClZr[376],Hg[377],P[378],Pd ₃ S ₂ Tl ₂ [379]
187	$Z_3 \times Z_3$	Br ₂ Ca ₃ Si[380],CHf[381],InNbS ₂ [382, 383],InNbSe ₂ [383, 384],NY[385]
189	$Z_3 \times Z_3$	AgAsCa[386, 387],AgCaP[387, 388] GeLiY[389]
191	$Z_6 \times Z_{12}$	B ₂ Ca[390],B ₂ Mg[391],B ₂ Ti[392],B ₂ Zr[393], Be ₁₂ Ti[394],Ga ₂ Sr[395],Pd ₅ Sr[396]
193	Z_{12}	Pb ₃ SZr ₅ [397]
194	Z_{12}	AlLi[398],AsHgK[399, 400], B ₂ W[401], Be[402],C[403],CdNa ₂ Sn[404], HgKSb[400, 405],SnTi ₂ [406]
221	$Z_4 \times Z_8$	CPbPd ₃ [407],CPt ₃ Sn[407],CaPd[408],Ca ₃ GeO[409, 410],Ca ₃ OPb[410, 411],HfPd ₃ [412],HgPt ₃ [413],NTi ₃ Tl[414] NY[415],PbPt ₃ [416],Pt ₃ Sn[417],Pt ₃ Zn[418]
225	Z_8	AsSc[419],Au ₂ InSc[420],Au ₂ InY[421],Ba[422],Bi ₃ [423],BiSc[424] CPt[425],Ca[426],PbPo[427, 428],SSn[429],SbSc[430],SeSn[431], SnTe[432, 433]
227	Z_4	Mo ₂ O ₇ Y ₂ [434],S ₂ Ti[435]

List 165 good topological Crystalline insulator

SG	Topological (semi-)metals
11	Br ₉ TeW[436], CBrHgNS[437], Li ₇ Sn ₃ [438], Mo ₂ S ₂ Sb[439]
51	AuCd[440], AuTi[441]
52	Ag ₂ BiO ₃ [442], Bi ₃ Sr ₂ [443]
55	Al ₃ Pd ₅ [444], Al ₃ Pt ₅ [445], BCl ₆ Sc ₄ [446], Bi ₉ Ca ₉ Cd ₄ [447], Bi ₉ Ca ₉ Zn ₄ [448], Bi ₉ Cd ₄ Sr ₉ [447], In ₅ S ₁₃ Y ₄ [449]
57	AlCaPd[450], BiK ₂ Sn[451]
58	C ₆₀ K[452]
59	Ag ₃ Sn[453]
60	F ₄ NaTi[454]
61	AgF ₂ [455]
62	AgAuP ₇ [456], AgF ₃ K[457], AlPt ₂ [458] Bi ₃ Ca ₅ [459], Bi ₃ Sr ₅ [460], Ca ₅ Sb ₃ [461] GeNb ₃ Te ₆ [462], GePd ₂ Y[463], N ₃ Nb ₂ [464], Nb ₃ Si[465] PdSbZr[466], SiTa ₃ Te ₆ [466]
63	AgCa[467], AuCa[468], BiZr[469], Ga ₃ PdSr[470], Ga ₅ Zr ₃ [471], GeSc[472], GeY[473], HfSb[474], K ₃ O ₄ Pd ₂ [475], K ₃ O ₄ Pt ₂ [476], K ₄ P ₃ [477, 478], N ₆ NaTa ₃ [479], PdY[480], SiY[481], Sr ₃ Tl ₅ [482]
64	AgCs ₂ F ₄ [483], Au ₁₀ Ca ₄ In ₃ [484], Bi[485, 486]
67	Ba ₉ In ₄ [487], In ₄ Pd ₁₇ Se ₄ [488], Pt ₁₁ Zr ₉ [489], Pt ₁₂ Si ₅ [490]
88	Al ₂₁ Pt ₈ [491], CsFO ₃ S[492], Ge ₈ Pd ₂₁ [493]
123	AgPPt ₅ [494], AlPPt ₅ [494], AsInPd ₅ [495], AsPd ₅ Tl[496] AsPt ₅ Tl[496], As ₃ BaPd ₂ [497], BaP ₃ Pd ₂ [498], CaPb[499] CdPd[500], CdPd ₅ Se[501], CdPt[502], Cd ₃ Zr[503] FKNb ₄ O ₅ [504], HgPd[505], HgPd ₅ Se[506], HgPt[507], Hg ₂ Pt[507] InPPd ₅ [508], InPPt ₅ [508], PPd ₅ Tl[508], PPt ₅ Tl[508] PdTi[509], Pd ₅ SeZn[510], SiSr[511], Sr[512]
127	AlSc ₂ Si ₂ [513], Au ₂ Ca ₂ Pb[514], Au ₂ InY ₂ [515], B ₂ Ta ₃ [516] B ₄ W[517], C ₂ B ₂ Y[518], Ga ₂ MgSc ₂ [519], Ca ₃ Hg ₂ [520] Ga ₂ Nb ₃ [521], Ga ₂ Ta ₃ [522], Ge ₂ Hf ₃ [523], Hg ₂ Sr ₃ [524, 525] InPd ₂ Y ₂ [526], In ₅ Ti ₂ [527], LiSi ₂ Y ₂ [528], PbPd ₂ Y ₂ [529]
129	AgMgSb[530], AsNbSi[531], AsSiTa[532], BaMgSi[533], BiKMg[534], GeTeZr[535], MoNTa[536]
130	AlMg ₄ Si ₃ [537], Se ₃ Tl ₅ [538]
131	OPd[539]
136	AlNb ₂ [540], AlTa ₂ [541], Bi ₂ MgO ₆ [542], O ₂ Pb[543, 544], O ₂ Pd[545], O ₂ Pt[546]
137	As ₂ Cd ₃ [547, 548], Bi ₂ Se ₃ [549]
139	Ag ₂ CaGe ₃ [550], Ag ₂ Ge ₂ Sr[550], Ag ₂ Si ₂ Sr[551], Ag ₂ Sn ₂ Sr[552], Al ₂ BaSi ₂ [553], Al ₂ Pb ₂ Sr[554], Al ₃ Nb[555] Al ₃ Ta[556], Al ₃ Zr[557], Al ₄ Ba[558], Au ₄ CaCd ₂ [559], BaBi ₄ Cl ₂ O ₆ [560], BaIn ₄ [561], Be ₁₂ Pd[562], Be ₁₂ Pt[562] CaGa ₄ [563], Ca ₁₁ Sb ₁₀ [564], Ga ₃ Nb[565, 566], Ga ₃ Ta[567], In ₄ Sr[568], Pt ₃ Sb[569], S ₂ Ti[570], Sb ₁₀ Sr ₁₁ [571] Si ₂ W[572], ZnZr ₂ [573]
140	AgCsP ₃ [574], AgP ₃ Rb[574], AgIn ₂ [575], AlZr ₂ [576], AuSe ₃ Tl ₄ [577], AuTl ₂ [578], Au ₃ Cd ₅ [579] Au ₄ In ₂ K[580], Au ₄ In ₂ Rb[580] BGe ₂ Nb ₅ [581], BMo ₂ [582], BTa ₂ [583], BTi ₂ [584], BW ₂ [585], Ba[586], BeTa ₂ [587], Bi ₃ In ₄ Pb[588] Ca ₅ Pt ₃ [589], Ca ₅ Si ₃ [590] Cs ₉ InO ₄ [591], GaHf ₂ [592], GaPt ₃ [593], GaZr ₂ [594], Ga ₅ Pd[595], GeHf ₂ [596], Ge ₃ Mo ₅ [597] Ge ₂ Nb ₅ [598], Ge ₃ W ₅ [599], Nb ₅ SiSn ₂ [600] Nb ₅ Si ₃ [601], Pb ₃ Sr ₅ [602], PdTl ₂ [603], SZr ₂ [604] Si ₃ W ₅ [605], Sr[606], Te ₃ Tl ₅ [607]
163	AlF ₆ LiPd[608], F ₆ GaLiPd[609], In ₂ Pt ₃ [610]
164	AlCl ₃ [611], CSC ₂ [612], CaHg ₂ [613], CaSb ₂ Zn ₂ [614], Cl ₂ Ti[615], HfTe ₂ [616, 617], Hg ₇ K ₂ [618] Hg ₇ Rb ₂ [618], OTi ₂ [619], Sb ₂ SrZn ₂ [620], Se ₂ Ti[621]
166	BaPb ₃ [622], Be ₁₇ Hf ₂ [623], Be ₁₇ Nb ₂ [624], Be ₁₇ Ta ₂ [623], GeTe[625, 626], Pb ₂ Pd ₃ S ₂ [627] Pb ₂ Pd ₃ Se ₂ [628], Po[629], SbSn[630] AgInSe ₂ [631], As ₃ NaZn ₄ [632], Bi ₄ Te ₃ [633], C ₃ Al ₈ N ₄ [634], Cd ₄ KP ₃ [635], GeP ₃ [636], In ₂ Se ₃ [637], Pd ₃ Se ₂ Tl ₂ [638], Ag ₄ Sb ₂ Sr[639]
167	Al ₅ Mo[640], Cs ₈ Ga ₁₁ [641], F ₃ Mo[642], F ₃ Pd[643], F ₃ Ti[644], In ₁₁ K ₈ [645] K ₈ Tl ₁₁ [646], Mo ₉ S ₁₁ Tl ₂ [647], NPd ₃ [648]

List 489 good topological Semimetal

176	AgCd ₃ F ₂₀ Hf ₃ [649], AgCd ₃ F ₂₀ Zr ₃ [649], AgMo ₃ Se ₃ [650], AsNb ₃ Te ₃ [651] CsMo ₃ S ₃ [652], CsMo ₃ Se ₃ [653], CsMo ₃ Te ₃ [654], InMo ₃ Se ₃ [655, 656] InMo ₃ Te ₃ [655, 656], KMo ₃ S ₃ [655], KMo ₃ Se ₃ [655, 656], KMo ₃ Te ₃ [656, 657], LiMo ₃ Se ₃ [658] Mo ₃ NaSe ₃ [655, 656], Mo ₃ NaTe ₃ [656, 658], Mo ₃ RbS ₃ [655], Mo ₃ RbSe ₃ [658, 659] Mo ₃ RbTe ₃ [660], Mo ₃ Se ₃ Tl[656, 658], Mo ₆ Se ₆ Tl ₂ [661], NbSe ₂ [662]
187	AgN[663], CLiNaO ₃ [664], CMo[665], CPt[666], CTa[666], CW[667] InS ₂ Ta[668], InSe ₂ Ta[669], MoP[670], NNb[671], NPd[672], NW[673] NZr[674], N ₂ Pt[675], N ₂ Ta[675], NbS[676], STa[677] STi[678], TeZr[679]
188	I ₃ LiSc[680]
189	AlHPt[681], AlPdY[682], AlPtZr[681], Al ₂ Hf ₆ Pt[683], AsPd ₂ [684] AuInY[685], GaPtZr[681], GePdSc[686], GePd ₂ [687], GePt ₂ [688], InPdY[689] InPtSc[690], InPtY[691], MgPdY[692], NTa[693], PdTiY[694], PdYZn[695] Pd ₂ Si[696], PtSb ₂ Zr ₆ [697], Pt ₂ Si[698]
190	AlHPt[699], AlPtZr[700], B ₄ Ga ₃ Pt ₉ [701], GaHPd[702], GaPtZr[703], Li ₂ Sb[704], PdScSn[705]
191	Au ₂ Ba[706], Au ₅ K[707], Au ₅ Rb[707], B ₂ Mo[708], BaGa ₂ [709], BaPd ₅ [710], Be ₂ Hf[711] Be ₅ Hf[712], Be ₅ Zr[713], CaGa ₂ [714], CaHg ₂ [715], Hf[716], Hf ₂ N ₃ Ta[717], Hg[718] LiNNa ₂ [719], NNa ₃ [720], NOTa[721], NTa[722], N ₃ TaTi ₂ [723], N ₃ TaZr ₂ [723], N ₄ Ta ₂ [724], S ₂ Ti[725]
193	AgIn ₃ Zr ₅ [726], AgPb ₃ Zr ₅ [727], AgSb ₃ Zr ₅ [728], AlSn ₃ Zr ₅ [729], Al ₃ Hf ₅ [730] Al ₃ Hf ₅ N[731], Al ₃ Ta ₅ [732], Al ₃ Zr ₅ [733], Al ₄ Zr ₅ [734] AsPb ₃ Zr ₅ [735], AsSb ₃ Zr ₅ [736], AsSn ₃ Zr ₅ [737], As ₃ Ca ₅ [738], As ₃ Sr ₅ [739] BNb ₅ Si ₃ [740], BSn ₃ Zr ₅ [741], Ba ₃ N[742], Ba ₅ Bi ₃ [743] Ba ₅ Sb ₃ [743], Bi ₃ Sr ₅ [744], CSb ₃ Zr ₅ [745], CSn ₃ Zr ₅ [746], Ca ₅ Sb ₃ [747], CdPb ₃ Zr ₅ [748] Cl ₃ Ti[749, 750], Cl ₃ Zr[750, 751], GaSn ₃ Zr ₅ [752], Ga ₃ Hf ₃ Nb ₂ [753] Ga ₃ Nb ₅ [754], Ga ₃ Sc ₅ [755], Ga ₃ Ta ₅ [756], Ga ₄ Nb ₅ [757], Ga ₄ Ti ₅ [758] Ga ₄ Zr ₅ [758], GePb ₃ Zr ₅ [759], Ge ₃ Mo ₅ [760], Ge ₃ Nb ₅ [761] Ge ₃ Sc ₅ [762], Ge ₃ Ta ₅ [763], Ge ₃ Y ₅ [764], HfI ₃ [765, 766], Hf ₅ Sb ₃ Zn[767], Hf ₅ Si ₃ [768] Hf ₅ Sn ₃ [769], Hf ₅ Sn ₄ [770], Hg ₃ Mg ₅ [771], I ₃ Nb[772] I ₃ Ti[773, 774], I ₃ Zr[775], InPb ₃ Zr ₅ [776], K ₃ Nb ₅ O ₂₁ [777] Mo ₅ Si ₃ [778], NSn ₃ Zr ₅ [779], N ₆ Nb ₅ [780], N ₆ Ta ₅ [781] Nb ₅ OPt ₃ [782], Nb ₅ PSi ₃ [783], Nb ₅ Si ₃ [784], PPb ₃ Zr ₅ [785] PSn ₃ Zr ₅ [786], P ₃ Ti ₅ [787], Pb ₃ SbZr ₅ [788], Pb ₃ Sc ₅ [789] Pb ₃ SeZr ₅ [788], Pb ₃ SiZr ₅ [788], Pb ₃ SnZr ₅ [788], Pb ₃ Y ₅ [790] SSb ₃ Zr ₅ [791], SSn ₃ Zr ₅ [792], Sb ₃ SiZr ₅ [793] Sb ₃ Sr ₅ [794], Sb ₄ Zr ₅ [795], SeSn ₃ Zr ₅ [796], SiSn ₃ Zr ₅ [796] Si ₃ Ta ₅ [797], Si ₃ Y ₅ [798], Sn ₄ Zr ₅ [799]

*Tang, Po, Vishwanath, Wan**, *Nature* **566**, 486 (2019)

Zhang, Jiang, Song, Huang, He, Fang, Weng, Fang, *Nature* **566**, 475 (2019)

Vergniory, Elcoro, Felser, Bernevig, Wang, **566**, 480 *Nature* (2019)

Symmetry indicator

➤ TaAs family

Weng, Fang, Fang, Bernevig, & Dai, PRX (2015)

Huang et al., Nature Commun. (2015)

➤ Magnetic

Conclusion

□ We propose a highly efficient method to explore topological materials

What one need to do is to look the expansion coefficient!

□ We propose numbers of new topological materials.

**Thank you very much for your
attention**

ccmp.nju.edu.cn

*Tang, Po, Vishwanath, **Wan***, Nature Physics **15**, 470 (2019)*

*Tang, Po, Vishwanath, **Wan***, Science Advances **5**, eaau8725 (2019)*

*Tang, Po, Vishwanath, **Wan***, Nature **566**, 486 (2019)*

当前的问题和可能的方案

Na₃Bi

Wang, Sun, Chen, Franchini, Xu, Weng, Dai, and Fang, PRB (2012).

Cd₃As₂

Wang, Weng, Wu, Dai, and Fang, PRB (2013)

SnTe

Hsieh, Lin, Liu, Duan, Bansil and Fu, Nat. Commun. (2012).

Wang, West, Liu, Li, Yan, Gu, Zhang, and Duan, PRB (2014).